1	Forest Practice Committee Cumulative Impacts Assessment Discussion
2	November 2, 2016
3	
4	912.9, 932.9, 952.9 Cumulative Impacts Assessment Checklist [All Districts]
5	
6	STATE OF CALIFORNIA BOARD OF FORESTRY AND FIRE PROTECTION
7	CUMULATIVE IMPACTS ASSESSMENT
8	(1) Do the assessment area(s) of resources that may be affected by the proposed
9	project contain any past Past Projects, present projects, or Reasonably Foreseeable
10	Probably Future Projects reasonably foreseeable probable future projects? Yes
11	No
12	If the answer is yes, identify the project(s) and affected resource subject(s).
13	(2) Are there any continuing, significant adverse impacts from past land use
14	activities that may add to the impacts of the proposed project? Yes No
15	If the answer is yes, identify the activities, describing their location, impacts and affected
16	resource subject(s).
17	(3) Will the proposed project, as presented, in combination any past Past Projects
18	present <u>projects</u> , and <u>Reasonably Foreseeable Probably Future Projects reasonably</u>
19	foreseeable probable future projects identified in items (1) and (2) above, have a
20	reasonable potential to cause or add to significant cumulative impacts in any of the
21	following resource subjects?
22	

	Yes	No	No reasonably
	after mitigation	after mitigation	potential
	(a)	(b)	significant effects
			impacts (c)
1. Watershed			
2. Soil			
Productivity			
3. Biological			
4. Recreation			
5. Visual			
6. Traffic			
7. <u>Greenhouse</u>			
Gases (GHG)			
8. Wildfire Risk			
and Hazard			
9. Other			

- a) "Yes <u>after mitigation"</u>, means that potential significant adverse cumulative impacts are left after application of the forest practice rules Forest Practice Rules and mitigations or alternatives proposed by the plan submitter.
- b) "No after mitigation" means that any potential for the proposed timber operation to cause or add to significant adverse cumulative impacts by itself or in combination with other projects has been reduced to insignificance or avoided by mitigation measures or alternatives proposed in the THP Plan and application of the forest practice rules Forest Practice Rules.

Yes	No	No reasonably
after mitigation	after mitigation	potential
(a)	(b)	significant effects
		impacts (c)

- c) "No reasonably potential significant cumulative effects impacts" means that the operations proposed under the THP Plan do not have a reasonable potential to join with the impacts of any other project to cause, add to, or constitute significant adverse cumulative impacts.
- (4) If column (a) is checked in (3) above describe why the expected impacts cannot be feasibly mitigated or avoided and what mitigation measures or alternatives were considered to reach this determination. If column (b) is checked in (3) above describe what mitigation measures have been selected which will substantially reduce or avoid reasonably potential significant cumulative impacts except for those mitigation measures or alternatives mandated by application of the rules of the Board rules.

- (5) Provide a brief description of the assessment area used for each resource subject.
- (6) List and briefly describe the individuals, organizations, and records consulted in the assessment of cumulative impacts for each resource subject. Records of the information used in the assessment shall be provided to the Director upon request.

BOARD OF FORESTRY <u>AND FIRE PROTECTION</u> TECHNICAL RULE ADDENDUM NO. 2

CUMULATIVE IMPACTS ASSESSMENT

Introduction

The purpose of this addendum is to guide-provide the minimum standards included by RPFs within the assessment of cumulative impacts assessment as required in 14 CCR §§ 898, 912.9, 932.9, 952.9 and 1034 that may occur as a result of proposed timber operations. This assessment shall include evaluation of both on-site and off-site interactions of proposed project activities with the impacts of past and reasonably foreseeable future projects.

In conducting an a cumulative impacts assessment, the RPF must shall distinguish between on-site impacts that are mitigated by application of the California Forest Practice Rules and the interactions of proposed activities (which may not be significant when considered alone) with impacts of Past Projects and Reasonably Foreseeable Probable Future Projectspast and reasonably foreseeable future projects pursuant to PRC § 15130(b)(1)(a).

Resource subjects to bethat shall be considered in the preparation of a the assessment of cumulative impacts cumulative impacts assessment are described in the Appendix.include watershed resources, soil productivity, biological resources, recreational resources, visual resources, vehicular traffic impacts, greenhouse gas impacts and wildfire.

The RPF preparing a THP <u>Plan</u> shall conduct an assessment based on information that is reasonably available <u>before prior to</u> submission of the THP <u>Plan</u>. RPFs <u>are</u>

expected to shall submit sufficient information to the Director to support their findings if significant issues are raised during the Department's review of the THP Plan.

Information used in the assessment of a cumulative impacts assessment may be supplemented during the THP Plan review period. Agencies participating in plan Plan review may provide input into the cumulative impacts assessment based upon their area of expertise. Agencies should shall support justify and support their recommendations with documentation.

The Department, as lead agency, shall make the final determination regarding the sufficiency of a cumulative impacts assessment sufficiency and the presence or absence of significant cumulative environmental impacts. This determination shall be based on a review of all sources of information provided and developed during review of the Timber Harvesting Plan.

Identification of Resource Areas Geographic Assessment Areas

The RPF shall establish and briefly describe the geographic assessment area within or surrounding the <u>plan-Plan</u> for each resource subject to be assessed and shall briefly explain the rationale for establishing the <u>resource-geographic assessment</u> area. This shall be a narrative description and shall be shown on a map <u>where when</u> a map adds clarity to the <u>assessment</u> area.

Identification of Information Sources

The RPF shall list and briefly describe the individuals, organizations, and records used relied upon as sources of information in the assessment of preparation cumulative impacts assessment, including references for listed records and the names, affiliations, addresses, and phone numbers of specific individuals contacted. Records of information

Comment [A1]: Staff Comment: Is this sentence required give the previous sentence.

Comment [A2]: Staff comment: What type of documentation "relevant" "contemporary" "empirical"

1	used in the preparation of a cumulative impacts assessment shall be provided to the		
2	Director upon request.		
3	Common sources of information for cumulative effects impacts assessment are		
4	identified below. Sources to be used will depend upon the complexity of individual		
5	situations and the amount of information available	from other plans. Sources not listed	
6	below may have to be consulted based on individ	ual circumstances. Not all sources of	
7	information need to be consulted for every THP P	<u>lan</u> .	
8	1. Consultation with Experts and Organizations:		
9	(a) County Planning Department;	(b) Biologists;	
10	(c) Geologists;	(d) Soil Scientists;	
11	(e) Hydrologists;	(f) Federal Agencies;	
12	(g) State Agencies;	(h) Public and private utilities.	
13	2. Records Examined:		
14	(a) Soil Maps;	(b) Geology Maps;	
15	(c) Aerial Photographs;	(d) Natural Diversity Data Base;	
16	(e) THP Plan Records;	(f) Special Environmental Reports;	
17	(g) Topographic Maps;	(h) Basin Plans;	
18	(i) Fire History Maps;		
19	(j) Relevant Federal Agency Docume	ents or Plans;	
20	(k) Relevant Watershed or Wildlife S	studies (published or unpublished);	
21	(I) Available Modeling Approaches		
22			
23	As provided in Section14 CCR § 898 of the rules	s, the RPF or supervised designee and	
24	the plan submitter must consult information source	es that are reasonably available.	

Past Projects and Reasonably Foreseeable Probable and Future Activities

2 Projects

1

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

- 3 Past Projects and Reasonably Foreseeable Probable Future Projects future projects
- 4 included in the cumulative impacts assessment shall be described as follows:
- 5 A. Identify and briefly describe the location of past and reasonably foreseeable probable
- 6 future projects Past Projects and Reasonably Foreseeable Probable Future Projects as
- defined in 14 CCR § 895.1 within described resource geographic assessment areas.
- 8 Include a map or maps and associated legend(s) clearly depicting the following
- 9 information:
 - 1. Township and Range numbers and Section lines.
 - 2. Boundary of the <u>pPlanning <u>wW</u>atershed(s) within which the plan area is located along with the CALWATER 2.2 identification number.</u>
 - 3. Location and boundaries of Past Projects, present projects and Reasonably

 Foreseeable Probable Future Projectspast, present and reasonably foreseeable

 probable future timber harvesting projects on land owned or controlled by the timberland
 owner of the proposed timber harvest within the pPlanning wWatershed(s) depicted in
 section (2) above. For purposes of this section, Past Projects past projects shall be
 limited to those projects submitted within ten years prior to submission of the THPPlan.
 - **4.** Silvicultural methods for each of the timber harvesting projects depicted in section (3) above. Each specific silvicultural method must be clearly delineated on the map(s), and associated THP Plan number referenced in the legend or an annotated list. In addition, shading, hatching, or labeling shall be used which clearly differentiates silvicultural methods into one of the four categories outlined in Table 1.
 - 5. A north arrow and scale bar (or scale text).
 - **6.** Source(s) of geographical information.

Comment [A3]: Strike or modify language to make congruent with 895.1.

The map scale shall be large enough to clearly represent one planning watershed per page or of a scale not less than 1:63,360. Planning watersheds with densely situated or overlapping harvest units, or those which are large or irregular in size, may require multiple maps to achieve clarity. Map(s) shall be reproducible on black & white copiers, and submitted on an 8½ x 11 page(s).

3 Table 1

Silvicultural Category	Silvicultural Method
Evenaged	Clearcutting, Seed Tree Seed Step, Seed Tree
Management	Removal Step, Shelterwood Preparatory Step,
14 CCR § 913.1 [933.1,	Shelterwood Seed Step, Shelterwood Removal Step
953.1]	
Unevenaged	Selection, Group Selection, Transition
Management	
14 CCR § 913.2 [933.2,	
953.2]	
Intermediate	Commercial Thinning, Sanitation-Salvage
Treatments	
14 CCR § 913.3 [933.3,	
953.3]	
Special	Special Treatment Area Prescriptions, Rehabilitation
Prescriptions and	of Understocked Area Prescription,
Other Management	Fuelbreak/Defensible Space, Southern Subdistrict
14 CCR § 913.4 [933.4,	Special Harvesting Method (14 CCR § 913.8),
953.4]	Variable Retention, Conversion

Alternative Prescriptions shall be put into the category within which the most nearly appropriate or feasible silvicultural method in the Forest Practice Rules is found pursuant to 14 CCR § 913.6 (b)(3)[933.6(b)(3), 953.6(b)(3)].



B. The RPF shall Identify identify and give the location and description of any known, continuing significant environmental problems effects caused by past Past projects

Projects as defined in 14 CCR § 895.1. The RPF who prepares the plan Plan, or their supervised designee, shall obtain information from plan submitters (timberland or timber owner), and from appropriate agencies, landowners, and individuals about past, and future land management activities and shall consider past experience, if any, in the assessment area related to past impacts and the impacts of the proposed operations, rates of recovery, and land uses. A poll of adjacent land owners is encouraged and may be required by the Director to determine such activities and significant adverse environmental problems on adjacent ownerships.

Comment [A4]: Staff Comment: "Geographic" "watershed" or other assessment area.

Comment [A5]: Not an enforceable standard. Enforceability needs to be addressed for this standard.

Appendix Technical Rule Addendum # 2

In evaluating cumulative impacts, the RPF shall consider the factors set forth herein.

A. Watershed Resources

Cumulative Watershed Effects (CWEs) occur within and near bodies of water or significant wet areas wet meadows or other wet areas, where individual impacts are combined to produce an effect that is greater than any of the individual impacts acting alone. Factors to consider in the evaluation of cumulative watershed impacts are listed below.

- 1. Impacts to watershed resources within the Watershed Assessment Area (WAA) shall be evaluated based on significant on-site and off-site cumulative effects on beneficial uses of water, as defined and listed in applicable Water Quality Control Plans.
- 2. Watershed effects produced by timber harvest and other activities may include one or more of the following:

- Sediment
- Water temperature
- Organic debris
- Chemical contamination
- Peak flow

- The following general guidelines shall be used considered when evaluating watershed impacts. The factors described are general and may not be appropriate for all situations. Actual measurements may be required if needed to evaluate significant environmental effects. The plan must comply with the quantitative or narrative waterquality objectives set forth in an applicable Water Quality Control Plan.
 - a. Sediment Effects. Sediment-induced CWEs occur when earth materials transported by surface or mass wasting erosion enter a stream or stream system at separate locations and are then combined at a downstream location to produce a change in water quality or channel condition. The eroded materials can originate from the same or different projects. Sediment is composed of both suspended and bedload material. Suspended sediment is usually the primary source of turbidity in forested watersheds, although suspended organic material also accounts for a proportion of the suspended load. Chronic turbidity can be an indicator of a cumulative watershed sediment effect when sources can be identified and linked to one or more projects. Both turbidity and suspended sediment concentrations are subject to extreme inherent variability from region to region, storm to storm, and from year to year, dependent upon underlying geology and precipitation.

Potentially adverse <u>sediment</u> changes are most likely to occur in the following locations and situations:

1	 Downstream areas of reduced low stream gradient where
2	sediment from a new source may be deposited in addition to sediment derived from
3	existing or other new sources.
4	- Immediately downstream from where sediment from a new
5	source is combined with sediment from other new or existing sources and the combined
6	amount of sediment exceeds the transport capacity of the stream.
7	- Any location where sediment from new sources in
8	combination with suspended sediment from existing or other new sources significantly
9	increases turbidity- reduces the survival of fish or other aquatic organisms- or otherwise
10	reduces the quality of waters used for domestic, agricultural, or other beneficial uses.
11	- Channels with relatively steep gradients which contain
12	accumulated sediment and debris that can be mobilized by sudden new sediment inputs,
13	such as debris flows, resulting in debris torrents and severe channel scouring.
14	Potentially significant adverse impacts of cumulative sediment inputs
15	may include:
16	 Increased treatment needs or reduced suitability for domestic,
17	municipal, industrial, or agricultural water use.
18	- Direct mortality of fish and other aquatic species.
19	- Impaired spawning and rearing habitat for salmonids or
20	otherwise Rreduced viability of aquatic organisms or disruption of aquatic habitats and
21	loss of stream productivity caused by filling of pools and plugging or burying streambed
22	gravel.
23	- Accelerated channel filling (aggradation) resulting in loss of
24	streamside vegetation and stream migration that can cause accelerated bank erosion.

1	- Accelerated channel filling (aggradation) resulting in increased
2	frequency and magnitude of overbank flooding.
3	- Accelerated filling of downstream reservoirs, navigable
4	channels, water diversion and transport facilities, estuaries, and harbors.
5	- Channel scouring by debris flows and torrents.
6	- Nuisance to or reduction in water related recreational
7	activities.
8	Situations where sediment production potential is greatest include:
9	- Sites with high or extreme erosion hazard ratings.
10	- Sites which are tractor logged on steep slopes.
11	- Unstable areas.
12	b. Water Temperature Effect. Water temperature related CWEs are
13	changes in water chemistry or biological properties caused by the combination of solar
14	warmed water from two or more locations (in contrast to an individual effect that results
15	from impacts along a single stream segment) where natural cover has been removed.
16	Cumulative changes in water temperature are most likely to occur in the following
17	situations:
18	- Where stream bottom materials are dark in color.
19	- Where water is shallow and has little underflow.
20	- Where removal of streamside canopy results in substantial,
21	additional solar exposure or increased contact with warm air at two or more locations
22	along a stream.
23	- Where removal of streamside canopy results in substantial,
24	additional solar exposure or increased contact with warm air at two or more streams that
25	are tributary to a larger stream.

1	- Where water temperature is near a biological threshold for
2	specific species.
3	Significant adverse impacts of cumulative temperature increases
4	include:
5	- Increases in the metabolic rate of aquatic species.
6	- Direct increases in metabolic rate and/or reduction of
7	dissolved oxygen levels, either of which can cause reduced vigor and death of sensitive
8	fish and other sensitive aquatic organisms.
9	- Increased growth rates of microorganisms that deplete
10	dissolved oxygen levels or increased disease potential for organisms.
11	- Stream biology shifts toward warmer water ecosystems.
12	c. Organic Debris Effects. CWEs produced by organic debris can
13	occur when logs, limbs, and other organic material are introduced into a stream or lake at
14	two or more locations. Decomposition of this debris, particularly the smaller sized and
15	less woody material, removes dissolved oxygen from the water and can cause impacts
16	similar to those resulting from increased water temperatures. Introduction of excessive
17	small organic debris can also increase water acidity.
18	Large organic debris is an important stabilizing agent that should be maintained in
19	small to medium size, steep gradient channels, but the sudden introduction of large,
20	unstable volumes of bigger debris (such as logs, chunks, and larger limbs produced
21	during a logging operation) can obstruct and divert streamflow against erodible banks,
22	block fish migration, and may cause debris torrents during periods of high flow.
23	Removing streamside vegetation can reduce the natural, annual inputs of litter to the

stream (after decomposition of logging-related litter). This can cause both a drop in food

supply, and resultant productivity, and a change in types of food available for organisms

24

that normally dominate the lower food chain of streams with an overhanging or adjacent forest canopy.

d. Chemical Contamination Effects. Potential sources of chemical CWEs include run-off from roads treated with oil or other dust-retarding materials, direct application or run-off from pesticide treatments, contamination by equipment fuels and oils, and the introduction of nutrients released during slash burning or wildfire from two or more locations.

e. Peak Flow Effects. CWEs can be caused by management induced peak flow increases in streams during storm events. are difficult to anticipate. Peak flow increases may result from management activities that reduce rainfall interception (i.e., evaporation) and vegetative water use (i.e., transpiration), or produce openings where snow can accumulate, (such as clear-cutting in-clearcuts and site preparation on roads and landings), or that change the timing of flows by producing more efficient runoff runoff (such as insloped roads). These While increases, if any, however, are likely to be small relative to pre-harvest natural peak flows, extensive canopy removal over a short period of time on a watershed scale can increase peak flow effect on streambank erosion, channel incision, and headward channel extension in erodible landscapes. from medium and large storms. Research to date on the effects of management activities on channel conditions indicates that channel changes during storm events are primarily the result of large sediment inputs. The timing and concentration of flows affecting lower order stream channel morphology can also be affected by the routing of runoff from roads, landings, and skid trails. Peak flow effects diminish with decreasing intensity of canopy removal, increasing time since harvest, and during larger flow recurrence intervals.

1

2

3

4

5

6 7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

3. Watercourse Condition. The watershed impacts of past upstream and 1 on-site projects are often reflected in the condition of stream channels on the project area. 2 3 Following is a list of channel characteristics and factors that may be used to describe current watershed conditions and to assist in the evaluation of potential project impacts: 4 ♦ Gravel Embedded - Spaces between stream gravel filled with sand 5 or finer sediments. Gravel are often in a tightly packed arrangement. 6 ♦ Pools Filled - Former pools or apparent pool areas filled with 7 sediments leaving few areas of deep or "quiet" water relative to stream flow or size. 8 ♦ Aggrading - Stream channels filled or filling with sediment that raises 9

♦ Aggrading - Stream channels filled or filling with sediment that raises the channel bottom elevation. Pools will be absent or greatly diminished and gravel may be embedded or covered by finer sediments. Streamside vegetation may be partially or completely buried, and the stream may be meandering or cutting into its banks above the level of the former streambed. Depositional areas in aggrading channels are often increasing in size and number.

10

11

12

13

14

15

16

17

18

19

20

21

22

23

♦ Bank Cutting - Can either be minor or severe and is indicated by areas of fresh, unvegetated soil or alluvium exposed along the stream banks, usually above the low-flow channel and often with a vertical or undercut face. Severe bank cutting is often associated with channels that are downcutting, which can lead to over-steepened banks, or aggrading, which can cause the channel to migrate against slopes that were previously above the high flow level of the stream.

♦ Bank Mass Wasting - Channels with landslides directly entering the stream system. Slide movement may be infrequent (single events) or frequent (continuing creep or periodic events). ♦ Downcutting - Incised stream channels with relatively clean, uncluttered beds cut below the level of former streamside vegetation and with eroded, often undercut or vertical, banks.

♦ Scoured - Stream channels that have been stripped of gravel and finer bed materials by large flow events or debris torrents. Streamside vegetation has often been swept away, and the channel has a raw, eroded appearance.

♦ Organic Debris - Debris in the watercourse can have either a positive or negative impact depending on the amount and stability of the material. Some stable organic debris present in the watercourse helps to form pools and retard sediment transport and downcutting in small to medium sized streams with relatively steep gradients. Large accumulations of organic debris can block fish passage, block or divert streamflow, or could be released as a debris flow.

♦ Stream-Side Vegetation - Stream-side vegetation and near-stream vegetation provide shade or cover to the stream, which may have an impact on water temperature, and provides root systems that stabilize streambanks and floodplains and filter sediment from flood flows.

♦ Recent Floods - A recent high flow event that would be considered unusual in the project area may have an impact on the current watercourse condition.

B. Soil Productivity

Cumulative soil productivity impacts occur when the effects of two or more activities, from the same or different projects, combine to produce a significant decrease in soil biomass production potential. These impacts most often occur on-site within the project boundary, and the relative severity of productivity losses for a given level of impact generally increases as site quality declines. The primary factors influencing soil productivity that can be affected by timber operations include:

♦ Organic matter loss.
♦ Soil compaction.

2 ♦ Surface soil loss. ♦ Growing space loss.

The following general guidelines may be used when evaluating soil productivity impacts.

1. Organic Matter Loss. Displacement or loss of organic matter can result in a long term loss of soil productivity. Soil surface litter and downed woody debris are the store-house of long term soil fertility, provide for soil moisture conservation, and support soil microorganisms that are critical in the nutrient cycling and uptake process. Much of the chemical and microbial activity of the forest nutrient cycle is concentrated in the narrow zone at the soil and litter interface.

Displacement of surface organic matter occurs as a result of skidding, mechanical site preparation, and other land disturbing timber operations. Actual loss of organic matter occurs as a result of burning or erosion. The effects of organic matter loss on soil productivity may be expressed in terms of the percentage displacement or loss as a result of all project activities.

2. Surface Soil Loss. The soil is the storehouse of current and future site fertility, and the majority of nutrients are held in the upper few inches of the soil profile. Topsoil displacement or loss can have an immediate effect on site productivity, although effects may not be obvious because of reduced brush competition and lack of side-by-side comparisons or until the new stand begins to fully occupy the available growing space.

Surface soil is primarily lost by erosion or by displacement into windrows, piles, or fills. Mass wasting is a special case of erosion with obvious extreme effects on site productivity. The impacts of surface soil loss may be evaluated by estimating the proportion of the project area affected and the depth of loss or displacement.

1	3. Soil Compaction. Compaction affe	ects site productivity through loss of
2	large soil pores that transmit air and water in the so	oil and by restricting root penetration.
3	The risk of compaction is associated with:	
4	- Depth of surface litter.	- Soil structure.
5	- Soil organic matter content.	- Presence and amount of coarse
6	fragments in the soil.	
7	- Soil texture.	- Soil moisture status.
0		

Compaction effects may be evaluated by considering the soil conditions, as listed above, at the time of harvesting activities and the proportion of the project area subjected to compacting forces.

4. Growing Space Loss. Forest growing space is lost to roads, landings, permanent skid trails, and other permanent or non-restored areas subjected to severe disturbance and compaction.

The effects of growing space loss may be evaluated by considering the overall pattern of roads, etc., relative to feasible silvicultural systems and yarding methods.

C. Biological Resources

Biological assessment areas will vary with the species being evaluated and its habitat. Factors to consider in the evaluation of cumulative biological impacts include:

1. Any known rare, threatened, or endangered species or sensitive species (as described in the Forest Practice Rules) that may be directly or indirectly affected by project activities. Significant cumulative effects on listed species may be expected from the results of activities over time which combine time which combines to have a substantial effect on the species or on the habitat of the species.

2. Any significant, known wildlife or fisheries resource concerns within the immediate project area and the biological assessment area (e.g. loss of oaks creating forage problems for a local deer herd, species requiring special elements, sensitive species, and significant natural areas). Significant cumulative effects may be expected where there is a substantial reduction in required habitat or the project will result in substantial interference with the movement of resident or migratory species.
The significance of cumulative impacts on non-listed species viability should be determined relative to the benefits to other non-listed species. For example, the manipulation of habitat results in conditions which discourage the presence of some species while encouraging the presence of others.

- 3. The aquatic and near-water habitat conditions on the THP Plan and immediate surrounding area. Habitat conditions of major concern are: Pools and riffles, Large woody material in the stream, Near-water vegetation. Much of the information needed to evaluate these factors is described in the preceding Watershed Resources section. A general discussion of their importance is given below:
- a. Pools and Riffles. Pools and riffles affect overall habitat quality and fish community structure. Streams with little structural complexity offer poor habitat for fish communities as a whole, even though the channel may be stable. Structural complexity is often lower in streams with low gradients, and filling of pools can reduce stream productivity.
- b. Large Woody Material. Large woody debris in the stream plays an important role in creating and maintaining habitat through the formation of pools. These pools comprise important feeding locations that provide maximum exposure to drifting food organisms in relatively quiet water. Removal of woody debris can reduce frequency and quality of pools.

1	c. Near-Water Vegetation. Near-water vegetation provides many
2	habitat benefits, including: shade, nutrients, vertical diversity, migration corridors, nesting
3	roosting, and escape. Recruitment of large woody material is also an important element
4	in maintaining habitat quality.

- **4.** The biological habitat condition of the THP Plan and immediate surrounding area. Significant factors to consider are:
- ♦ Snags/den trees
 ♦ Hardwood cover
- 8 O Downed, large woody debris Late seral (mature) forest characteristics.
- 9 ♦ Multistory canopy ♦ Late seral habitat continuity
- 10 ♦ Road density

The following general guidelines may be used when evaluating biological habitat. The factors described are general and may not be appropriate for all situations. The THP Plan preparer must also be alert to the need to consider factors which are not listed below. Each set of ground conditions are unique and the analysis conducted must reflect those conditions.

- a. Snags/Den/Nest Trees: Snags, den trees, nest trees and their recruitment are required elements in the overall habitat needs of more than 160 wildlife species. Many of these species play a vital role in maintaining the overall health of timberlands. Snags of greatest value are >16" DBH and 20 ft. in height. The degree of snag recruitment over time should be considered. Den trees are partially live trees with elements of decay which provide wildlife habitat. Nest trees have importance to birds classified as a sensitive species.
- b. Downed large, woody debris: Large downed logs (particularly conifers)
 in the upland and near-water environment in all stages of decomposition provide an
 important habitat for many wildlife species. Large woody debris of greatest value consists

- of downed logs >16" diameter at the large end and >20 feet in length.
- 2 **c. Multistory canopy**: Upland multistoried canopies have a marked
- 3 influence on the diversity and density of wildlife species utilizing the area. More
 - productive timberland is generally of greater value and timber site capability should be
- 5 considered as a factor in an assessment. The amount of upland multistoried canopy may
- 6 be evaluated by estimating the percent of the stand composed of two or more tree layers
- 7 on an average per acre basis.
- 8 Near-water multistoried canopies in riparian zones that include conifer and hardwood
- 9 tree species provide an important element of structural diversity to the habitat
- requirements of wildlife. Near-water multistoried canopy may be evaluated by estimating
 - the percentage of ground covered by one or more vegetative canopy strata, with more
- emphasis placed on shrub species along Class III and IV streams (14 CCR §§ 916.5,
- 13 936.5, or 956.5).

11

16

17

18

19

20

23

- 14 d. Road Density: Frequently traveled permanent and secondary roads
- have a significant influence on wildlife use of otherwise suitable habitat. Large declines in
 - deer and bear use of areas adjacent to open roads are frequently noted. Road density
 - influence on large mammal habitat may be evaluated by estimating the miles of open
 - permanent and temporary roads, on a per-section basis, that receive some level of
 - maintenance and are open to the public. This assessment should also account for the
 - effects of vegetation screening and the relative importance of an area to wildlife on a
- seasonal basis (e.g. winter range).
- 22 **e. Hardwood Cover**: Hardwoods provide an important element of habitat
 - diversity in the coniferous forest and are utilized as a source of food and/or cover by a
 - large proportion of the state's bird and mammal species. Productivity of deer and other
- 25 species has been directly related to mast crops. Hardwood cover can be estimated using

the basal area per acre provided by hardwoods of all species.

[Northern and Southern only]: Post-harvest deciduous oak retention for the maintenance of habitats for mule deer and other hardwood-associated wildlife shall be guided by the Joint Policy on Hardwoods between the California Board of Forestry and California Fish and Game Commission (5/9/94). To sustain wildlife, a diversity of stand structural and seral conditions, and tree size and age classes of deciduous oaks should be retained in proportions that are ecologically sustainable. Regeneration and recruitment of young deciduous oaks should be sufficient over time to replace mortality of older trees. Deciduous oaks should be present in sufficient quality and quantity, and in appropriate locations to provide functional habitat elements for hardwood-associated wildlife.

f. Late Seral (Mature) Forest Characteristics:

and their structural characteristics provides characteristics provide a basis from which to begin an assessment of the influence of management on associated wildlife. These characteristics include large trees as part of a multilayered canopy, large decadent trees and the presence of a large numbers of snags and downed logs, all of which that contribute to an increased level of stand decadence and complexity. Late seral stage forest amount may be evaluated by estimating the percentage of the land base within the project and the biological assessment area occupied by areas conforming to the following definitions:

Forests not previously harvested should be at least 80 acres in size to maintain the effects of edge. This acreage is variable based on the degree of similarity in surrounding areas. The area should include a multi-layered canopy, two or more tree species with

several large coniferous trees per acre (smaller subdominant trees may be either conifers

- or hardwoods), large conifer snags, and an abundance of large woody debris.
- 2 Previously harvested forests are in many possible stages of succession and may
- 3 include remnant patches of late seral stage which generally conform to the definition of
- 4 unharvested forests but do not meet the acreage criteria.
- g. Late Seral Habitat Continuity: Projects containing areas meeting the
- 6 definitions for late seral stage characteristics must be evaluated for late seral habitat
- 7 continuity. The fragmentation and resultant isolation of late seral habitat types is one of
- 8 the most significant factors influencing the sustainability of wildlife populations not
- 9 adapted to edge environments.
- This fragmentation may be evaluated by estimating the amount of the on-site number of
- acres within both the project area, and as wells as the biological assessment area
- occupied by portions of or entire late seral stands greater than at least 80 acres in size
- 13 (considering the mitigating influence of adjacent and similar habitat, if applicable) and less
- than one mile apart or connected by a corridor of similar habitat.
- h. Special Habitat Elements: The loss of a key habitat element may have
- 16 a profound effect on a species even though the habitat is otherwise suitable. Each
- species may have several key limiting factors to consider. For example, a special need
 - for some large raptors is large decadent trees/snags with broken tops or other features.
- 19 Deer may have habitat with adequate food and cover to support a healthy population size
 - and composition but dependent on a few critical meadows suitable for fawning success.
 - These and other key elements may need special protection.

D. Recreational Resources RECREATIONAL RESOURCES

- 23 The recreational assessment area is generally the area that includes the logging area
- 24 plus 300 feet.

18

20

21

22.

To assess recreational cumulative impacts:

- Identify the recreational activities involving significant numbers of people in and within 300 ft. of logging area (e.g., fishing, hunting, hiking, picnicking, camping).
- Identify any recreational Special Treatment Areas described in the Board rules on the plan area or contiguous to the area.

E. Visual Resources VISUAL RESOURCES

1

2

4

5

9

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

- The visual assessment area is generally the logging area that is readily visible to significant numbers of people who are no further than three miles from the timber operation. To assess visual cumulative effects:
 - Identify any Special Treatment Areas designated as such by the Board because of their visual values.
 - 2. Determine how far the proposed timber operation is from the nearest point that significant numbers of people can view the timber operation. At distances of greater than 3 miles from viewing points activities are not easily discernible and will be less significant.
 - **3.** Identify the manner in which the public identified in 1 and 2 above will view the proposed timber operation (from a vehicle on a public road, from a stationary public viewing point or from a pedestrian pathway).

F. Vehicular Traffic Impacts VEHICULAR TRAFFIC IMPACTS:

- The traffic assessment area involves the first roads not part of the logging area on which logging traffic must travel. To assess traffic cumulative effects:
- Identify whether any publicly owned roads will be used for the transport of wood products.
- Identify any public roads that have not been used recently for the transport of wood products and will be used to transport wood products from the

3. Identify any public roads that have existing traffic or maintenance 2 3 problems. 4. Identify how the logging vehicles used in the timber operation will change 4 the amount of traffic on public roads, especially during heavy traffic conditions. 5 6 7 G. Greenhouse Gas (GHG) Impacts Forest management effects sequestration and emission rates of forests to the extent 9 10 management activities affect forest inventory, growth, yield, and mortality. Timber operations and subsequent production of wood products can result in both the emission 11 and storage of GHGs. 12 13 Any one or a combination of the following options can be used to assess the potential for 14 significant cumulative GHG effects: 15 1. Incorporation by reference, or tiering from, a programmatic assessment that was 16 certified by the Board, CAL FIRE, or other State Agency, which analyzes the net 17 effects of GHG associated with forest management activities. 18 19 2. Application of a model or methodology quantifying an estimate of greenhouse gas 20 emissions resulting from the project. The model or methodology should at 21 minimum consider the following: 22 a. Inventory, growth, and harvest over a specified planning horizon 23 b. Projected forest sequestration over the planning horizon 24

proposed timber harvest.

25

c. Timber operation related emissions originating from logging equipment and

2	d. GHG emissions and storage associated with the production and life cycle of
3	manufatured wood products
4	
5	3. A qualitative analysis describing the extent to which the project in combination with
6	Past Project and Reasonably Foreseeable Probable Future Projects may increase
7	or reduce GHG emissions compared to the existing environmental setting. Such
8	analysis should disclose if a known threshold of significance for the project type
9	has been identified by the Board, CAL FIRE or State Agency, and if so, if the
10	project's emissions in combination with other forestry projects are anticipated to
11	exceed this threshold.
12	
13	H. Wildfire
14	Modifications to fuel loading through timber harvest activities may affect wildfire risk, hazard
15	and behavior. The alteration of these aspects of wildfire may negatively or positively affect
16	communities and/or infrastructure within proximity to the proposed project, particulary in the
17	wildland urban interface.
18	
19	Factors to consider that affect wildfire risk, hazard, and behavior include:
20	
	♦ Fire hazard severity rating ♦ Post-harvest logging area fuel
	<u>loading conditions</u>
	♦ Topography ♦ Public/private roads
	♦ Residential/commercial structures ♦ Historic monuments
	♦ Local/state/federal parks ♦ Domestic water sources

transportation of logs to manufacturing facility

♦ Schools and hospitals